



International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR)

ISSN:2347-6567

IJAMSCR | Volume 4 | Issue 1 | Jan – Mar - 2016
www.ijamscr.com

Research article

Medical research

Cervical Spondylosis-A Review

Suddhasheel Roy^{*1}, Vishwambhar Singh²

¹Department of ENT, Institute of medical sciences, Banaras Hindu University Varanasi, Uttar Pradesh, 221005, India

²Department of ENT, Institute of medical sciences, Banaras Hindu University Varanasi, Uttar Pradesh, 221005, India

*Corresponding author: Suddhasheel Roy

Email: sr3nov@gmail.com

ABSTRACT

Cervical pain is a very common complain encountered in ENT outdoors. Cervical spondylosis is one of the leading causes presenting with cervical pain. The different possible presentations, evaluation, pathophysiology and treatment of this disease are discussed in brief.

Keywords: Spondylosis, Cervical.

INTRODUCTION

Degenerative changes in amphithrodial joints and zygoapophyseal joints of cervical spine leads to an array of symptoms. This disease is named cervical spondylosis or cervical osteoarthritis¹. A lot of patients present in ENT outpatient department with various complaints pertaining to the disease. It affects 90% of population greater than 65 years of age². In this article we review the presentation, evaluation, pathophysiology and management of the disease briefly from an ENT perspective.

Presentation

Most cases are asymptomatic³. They are usually presented as cervical pain, radiculopathy or myelopathy². Cervical pain is mainly carried by supply of sinuvertebral nerve and posterior ramus branches^{3,4}. Radiculopathy is due to compression of nerve roots. It may be sensory, motor or autonomic.

Sensory involvement manifests as pain, paresthesia or hyperesthesia. Pain can be in neck, upper limbs, scapular region, head, in both arms upto fingers. Pain can be neuralgic which is sharp shooting in nature or myalgic, a dull pain in the corresponding myotome. Paresthetia follows the dermatome of the cervical nerve involved.

Myelopathy is due to compression of long tracts⁵. Motor symptoms range from stiffness, paraparesis to foot drop. Initial symptoms are upper motor neuron type while later symptoms are mostly lower motor neuron type⁶⁻¹¹. Sensory symptoms range from headache, rotator vertigo, numbness, myoclonic jerks to hiccups¹.

Evaluation

Physical examination shows limited and painful rotation, lateral flexion and extension of neck. Palpation of cervical vertebrae may elicit tenderness and irregular bony projections. Spasm of

paraspinal muscles is a very common finding. Radiological investigation is the mainstay in suspected cases. Static and flexion-extension x-rays in antero-posterior and lateral views are the initial investigation. X-rays may show loss of intervertebral space height, kyphotic changes, osteophytes, facet arthropathy, and fusion of adjacent vertebrae^{6, 7}. CT scans are used for the evaluation of spinal canal's shape and size, transverse foramina and joint spaces^{2, 8, 9}. While MRI gives the better picture of the spinal canal dimensions, the spinal cord, intervertebral discs, and cervical soft tissues^{2,10-11}.

Pathophysiology

Three consequential general events were described by Sokoloff¹. Initially there is a degeneration of the articular cartilage leading to denudation of the joint. Fibrillation of articular cartilage and remodeling of the periarticular structures including bone follows. The stiffness of the subchondral bone increases which leads secondarily to changes in the cartilage. Mechanical factors involved in the pathogenesis of cervical spondylosis have been categorized into two groups as static and dynamic factors¹⁴ (Table 1).

Table 1: Mechanical factors in pathogenesis of spondylosis.

Static	Congenital canal stenosis, disc herniation, degenerative osteophytic growth, hypertrophy of the ligamentum flavum, calcification of PLL.
Dynamic	Abnormal forces on the spinal column and cord during movement.

The first event in the pathogenesis is decreased in concentration of proteoglycan of hyaline cartilage^{13, 15}. Which results in shrinkage and loss of elasticity of nucleus pulposus and loss of disc height and loss of cervical lordosis. Loss of disc space is more pronounced in the ventral part^{15, 16}. This in turn results in the additional pressure on dorsal part and ultimately kyphotic changes¹⁵. These lead to the reduction in sagittal spinal canal diameter which is the main pathophysiological change³.

The other mechanism of reduction of spinal canal diameter is disc herniation. The degenerated disc usually herniates through the dorsal weak annulus fibrosis^{17, 18}. Reactive bone formation or osteophytes projection into the spinal canal is the third contributing factor. This bone formation starts in the bare bone area where the posterior longitudinal ligament has peeled off the vertebral body edge^{18, 19}. Mostly osteophytic formation is prominent in the area of C5-C6 and C6-C7 where cervical movements are maximum¹⁷. A fourth factor which often results in the decreased diameter

is the congenital spinal canal stenosis⁶. This magnifies the effect of the degenerative changes and leads to early onset myelopathy.

Spinal cord compressions may be due to osteophytes or hypertrophied ligamentum flavum^{14, 19}. Vascular compression symptoms may also surface due to compressions of anterior and posterior spinal arteries and other local vessels^{6, 15}.

Treatment

The disease is usually progressive in nature and the first intervention should be educating the patient about the natural course of the disease. Physical therapy like heat, ultrasound, heat pads, infrared lamps can be used to reduce the pain¹. Analgesics, anti-inflammatories and muscle relaxants can be used depending on the intensity of pain. Stretching and range-of-motion exercises including cervical, thoracic and lumbar areas should be advised. In cases of patients with complications such as radiculopathy, myelopathy, vascular and compressive symptoms neurologic and orthopaedic consultations should be done urgently.

REFERENCES

- [1]. John H.Bland. Disorders of the Cervical Spine. 2nd edition. W.B. Saunders Company: 2nd Edition.
- [2]. Jeffrey Mullin, Daniel Shedid, Edward Benzel. Overview of Cervical Spondylosis Pathophysiology and Biomechanics. WScJ 2: 89-97, 2011
- [3]. Morishita Y, Falakassa J, Naito M, Hymanson HJ, Taghavi C, Wang JC. The Kinematic Relationships of the Upper Cervical Spine Spine 34, 2642–2645, 2009

- [4]. Garfin SR: Cervical Degenerative Disorders: Etiology, Presentation, and imaging Studies. Instr Course Lect 49:335-338, 2000
- [5]. Harrop J, Hanna A, Silva M, Sharan A. Neurological Manifestations of Cervical Spondylosis: An Overview of Signs, Symptoms, and Pathophysiology. Neurosurgery 60:S1-14–S1-20, 2007
- [6]. Asgari S: Cervical Spondylotic Myelopathy, in Palmer JD (ed): Neurosurgery '96: Manuel of Neurosurgery. New York, Churchill Livingstone, 1996 pp750-754
- [7]. Crandall PH, Batzdorf U: Cervical Spondylotic Myelopathy. J Neurosurg 25:57-66, 1966
- [8]. Alker G: Neuroradiology of Cervical Spondylotic Myelopathy. Spine 13:850-853, 1988
- [9]. Ferguson RJ, Caplan LR: Cervical Spondylotic Myelopathy. Neurol Clin 3:373-382, 1985
- [10]. Fager CA: Results of Adequate Posterior Decompression in the Relief of Spondylotic Cervical Myelopathy. J Neurosurg 38:684-692, 1973
- [11]. McCormick WE, Steinmetz MP, Benzel EC: Cervical Spondylotic Myelopathy: Make the Difficult Diagnosis, then Refer for Surgery. Cleve Clin J Med 70:899-904, 2003
- [12]. Patten J: Neurological Differential Diagnosis. London, Springer-Verlag, 1977, pp 139-171
- [13]. Sokoloff L: The remodeling of articular cartilage. In Schattesskirchner M(ed):Rheumatology, vol 7. Basel:S karger, 1982:11
- [14]. White AA 3rd, Panjabi MM: Biomechanical Consideration in the Surgical Management of Cervical Spondylotic Myelopathy. Spine13:856-860, 1988
- [15]. Benzel EC: Biomechanics of Spine Stabilization. Rolling Meadows,American Association of Neurological Surgeons Publications,2001
- [16]. Emery SE: Cervical Spondylotic Myelopathy: Diagnosis and Treatment. J Am Acad Orthop Surg 9:376-388, 2001
- [17]. Garfin SR: Cervical Degenerative Disorders: Etiology, Presentation, and imaging Studies. Instr Course Lect 49:335-338, 2000
- [18]. Verbiest H: Chapter 23. The Management of Cervical Spondylosis. Clin Neurosurg 20: 62-294, 1973
- [19]. Schneider RC, Cherry G, Pantek H: The Syndrome of Acute Central Cervical Spinal Cord Injury; with Special reference to the Mechanisms Involved in Hyperextension Injuries of Cervical Spine. J Neurosurg 28:565-581, 1968

How to cite this article: Suddhasheel Roy, Vishwambhar Singh, Cervical Spondylosis-A Review. Int J of Allied Med Sci and Clin Res 2016; 4(1): 148-150.

Source of Support: Nil. **Conflict of Interest:** None declared.