



## International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR)

ISSN:2347-6567

IJAMSCR /Volume 4 | Issue 3 | July - Sep - 2016  
www.ijamscr.com

Research article

Pharmacology research

### *In-vitro* anthelmintic activity of methanolic leaf extract of *Abrus precatorius* Linn. (fabaceae)

S. Selvadurai\*, M.D. Dhana Raju, A. Venkateswara Rao

Department of Pharmacognosy, GIET School of Pharmacy, Rajahmundry, AP-533296

\*Corresponding author: Selvadurai S

Email id: sasdurai85@gmail.com

#### ABSTRACT

The present study was aimed to investigation of the anthelmintic potential of crude methanolic extract of the leaves of *Abrus precatorius* Linn. On Indian earth-worm "Pheretima posthum" and "Tubifex tubifex". Three concentrations (5, 10, 15 mg/ml) of methanolic extract were studied in activity which involved the determination of time of paralysis (vermifuge) and time of death (vermicidal) of the worms. Piperazine citrate (10 mg/ml) was used as reference standard drug whereas distilled water as control. Determination of paralysis time and death time of the worms were recorded. The result shows that methanolic extract possesses vermifuge activity and found to be effective as an anthelmintic. Therefore, the anthelmintic activity of the methanolic extract of *Abrus precatorius* Linn. Leaves have been reported.

**Keywords:** *Abrus Precatorius* Linn. Anthelmintic Activity, Piperazine Citrate, Methanolic Extract, Pheretima Posthum, Tubifex Tubifex.

#### INTRODUCTION

##### Scientific Classification

Division	- Angiospermae
Class	- Magnoliopsida
Order	- Fabales
Family	- Fabaceae
Genus	- <i>Abrus</i>
Species	- <i>Precatorius</i>

##### Macroscopy

Synonym	- <i>Rosary pea, john crow bead</i>
Biological Source	- <i>Abrus precatorius</i> Linn
Family	- Fabaceae

##### Vernacular Names

English	- Crab's eye, Jequirity
Hindi	- Rati, Gaunchi
Kannadam	- Gulaganji

Malayalam	- Kunni, Gundumani
Tamil	- Gundumani, kunthamani
Telugu	- Guruvinda

Parasitic infection including Helminthiasis is a critical serious problem in the tropical regions including the Asian countries which affects more than two billions of people worldwide. Helminths produce serious problem in human and other animals around the world specifically to the third world countries [5]. Different type of helminths infects the human and animals out of which intestinal round worms (*Ascaridia* sp.) are most common. The helminths parasites mainly subsist in human body in intestinal tract, but they are also found in tissue, as their larvae migrate towards them [9]. Despite this prevalence of parasitic infections, the research on the anthelmintic drug is sparse. According to the WHO, only a few drugs

are used in treatment of helminthes in humans. Anthelmintics from natural sources could play a key role in the treatment of these parasite infections. In view of this, attempts have been made to study the anthelmintic activity of traditional medicinal plants. For our studies, we used *Abrus precatorius* Linn. that belong to the Fabaceae family *Abrus precatorius* Linn., a highly medicinal plant, and commonly known as Indian Liquorice in English, *Gunchi*, *Ratti* in Hindi; Kannada-*Gulganji*; Malayalam-*Kumi*; Sanskrit-*Rektika*, *Gunja* and in Tamil as *Kuntumani*. on open lands. Its aerial parts are used for treating certain health problems like leucorrhoea, gonorrhoea, diarrhoea and dysentery. The roots, leaves and seeds are used for medicinal purpose. The roots and leaves are astringent, sweet, emetic, diuretic, anthelmintic and alexeteric. They are useful in cough pharyngodynia, pectoralgia, inflammation, strangury and vitiated condition of *vata*. The seeds are acrid, bitter, astringent, purgative, toxic abortifacient, aphrodisiac and trichogenous; seeds paste applied locally for sciatica, stiffness of shoulder joints, paralysis, leucoderma, ulcers, skin disease, wounds, alopecia, asthma, tubercular glands, stomatitis, hyperdipsia and fever [4,8].

## MATERIALS AND METHODS

### Collection of plant material

The plant material *Abrus precatorius* Linn leaves were collected from local area of Rajahmundry, East Godavari DT, AP. in the month of April 2015 and authenticated by the renowned botanist Prof. V. B. Sunitha, Dept. of Botany, Government Arts College, Rajahmundry. A voucher specimen (GCR/01/2015) was deposited in the college herbarium.

### Drugs and Chemicals

Piperazine citrate (Glaxo Smithkline), methanol, Distilled water were used during the experimental protocol. All the chemicals used are laboratory and analytical grade.

### Preparation of Plant Extract [6]

The fresh leaves were dried under shade. The dried leaves were cut into small pieces and powdered in a hand mill. Then 300 gm of coarse powder was extracted with 900 ml of extra pure Methanol successively in a Soxhlet extractor repeatedly for 48 hours. The extract was dried by solvent evaporation in a thermostat water bath at

50-60 °C temperature and the extract was kept in dessicator for the experiment.

### Preliminary Phyto-chemical Investigation [7]

The percentage yield of Methanol extract obtained was calculated and the extracts were subjected to number of proximate qualitative phyto-chemical analysis. These procedures are already reported by number of workers and used without any modification.

### Selection of Experimental Model [3]

Indian adult earthworms *Pheretima posthuma* and *Tubifex tubifex* were used to carry out the experiment. The earthworms were collected from the local supplier. Worms were washed with normal saline to remove all faecal matter. The earthworms of 7-12 cm in length and 0.4-0.5 cm in width were used for all the experimental protocol. Ready availability, anatomical and physiological resemblance of *Pheretima posthuma* and *Tubifex tubifex* made it to be used initially for in-vitro evaluation of anthelmintic activity.

### Experimental Design [1, 2, 8]

The anthelmintic activity was carried out as described by Ajaiyeoba [2], with minor modifications. The Indian earthworm the assay was performed on adult Indian earthworm *Pheretima posthuma*, due to its anatomical and physiological resemblance with the intestinal roundworm parasites of human beings. *Pheretima posthuma* worms are easily available and used as a suitable model for screening of anthelmintic drug. The assay was also performed on the aquarium worm, *Tubifex tubifex*, because they belong to same group of Annelida. Briefly, 20 ml formulations containing three different concentrations, of crude methanolic extract of leaf (5, 10 and 15 mg/ml in double distilled water) were prepared and six earthworms (same size) and approximately 5 g of *Tubifex* worms were placed in it. Both the test solution and standard drug solution were freshly prepared and 'time for paralysis' was noted when no movement of any sort could be observed except when the worms were vigorously shaken. The 'time for death' of worms was recorded after ascertaining that the worms neither moved when shaken vigorously nor when dipped in warm water at 50°. A maximum time period of 120 min was ascertained for the paralyzing as well as death time of *Pheretima posthuma* and *Tubifex tubifex* worms. Piperazine citrate (10 mg/ml) was used as reference

standard with distilled water as the vehicle control. All experiments were repeated thrice.

## RESULTS AND DISCUSSION

Preliminary phytochemical screening of crude extract of the leaves of *Abrus precatorius* L, revealed the presence of tannins, steroids, flavonoids, saponin, terpenoids etc. The presence of phytoconstituents is reported in table 1. From the observations made, a dose dependent paralytic effect much earlier and the time of death was observed table - 2. Although, methanolic extract showed anthelmintic activity in a dose-dependent manner but the methanolic extract appeared to be more effective for both types of worms. Evaluation of anthelmintic activity was compared with reference standard piperazine citrate. The methanolic extract of the leaves of *abrus precatorius* l, caused paralysis at 22.33 min. and time of death at 45.00 min. for *Pheretima posthuma* and 14.66 min as paralysis time and

20.66 min as death time for *Tubifex tubifex* worms respectively. The reference drug piperazine citrate showed the time of paralysis and time of death as 25.00 and 64.00 min, respectively. Considering the methanolic extract of leaf showed comparable activity, it would be important to identify the key phytoconstituents. The mean and SEM were analyzed statistically by ANOVA followed by Dunnett's test,  $P < 0.05$  being considered as significant.

## CONCLUSION

The *Abrus precatorius* L. leaf extracts has showed significant anthelmintic activity at all the tested doses when compared to control as vermifuge and vermicidal while highest activity exhibited by the higher conc. (15 mg/ml) which assures the ethno-medicinal claim. Hence, we can think about this herb as alternate source of anthelmintic drugs and also can generate new active lead for suitable anthelmintic drug.



Fig – 1 MACROSCOPY

Table – 1 Data Showing the Preliminary Phytochemical Screening of *Abrus precatorius*. L

Phytoconstituents	Petroleum ether extract	Chloroform extract	Acetone extract	Methanolic extract	Aqueous extract
Alkaloids	(+)	(+)	(+)	(+)	(-)
Carbohydrates	(-)	(+)	(+)	(+)	(+)
Glycosides	(-)	(-)	(-)	(+)	(+)
Flavonoids	(-)	(-)	(-)	(+)	(+)
Phytosterols	(-)	(-)	(-)	(-)	(-)
Fixed oils and Fats	(-)	(-)	(-)	(-)	(-)
Saponins	(-)	(-)	(-)	(-)	(+)
Phenolic compounds and Tannins	(+)	(-)	(+)	(+)	(+)
Lignins	(-)	(-)	(-)	(-)	(-)
Proteins and Free Amino acids	(-)	(-)	(-)	(-)	(+)
Gums and Mucilage	(-)	(-)	(-)	(-)	(-)

Table – 2 Anthelmintic activity of methanolic extract of *Abrus precatorius* L.

Groups	Conc mg/ml	<i>Pheretima posthuma</i>		<i>Tubifex tubifex</i>	
		Paralyzing time (min)	Death time (min)	Paralyzing time (min)	Death time (min)
Distilled water	-	-	-	-	-
Leaf extract (Methanolic)	5	60.66 ± 0.67*	80.66 ± 0.67*	63.00 ± 2.08*	75.33 ± 1.45*
	10	36.32 ± 0.88*	66.33 ± 0.88*	31.33 ± 0.67*	39.33 ± 0.88*
	15	22.33 ± 0.88**	45.00 ± 0.58*	14.66 ± 0.88*	20.66 ± 1.33*
Piperazine citrate	10	30.6 ± 0.06*	51 ± 0.006*	25.00 ± 1.16*	64 ± 0.88*
	15	20.13 ± 0.98*	40.19 ± 0.15*	19.01 ± 1.11*	50.07 ± 1.19*

Results are expressed as mean ± sem N=3; \*P<0.05 as compared to standards.

## REFERENCES

- [1]. Ajaiyeoba EO, Onocha PA and Olarenwaju OT In vitro anthelmintic properties of *Buchholzia coriacea* and *Gynandropsis gynandra* extract. *Pharm Bio* 39, 2001, 217- 220
- [2]. Chandan HS, Tapas AR, Sakarkar DM Anthelmintic activity of extracts of *Coriandrum sativum* Linn. in Indian earthworm. *Int. J Phytomedicine* 3, 2011, 36-40.
- [3]. Chatterjee KD Parasitology, Protozoology and Helminthology. Calcutta: Guha Ray Sree Saraswaty Pres Ltd.; 1967, 168-169.
- [4]. Dr. Ravindra Fernando; National Poisons Information Centre, Written a review about *Abrus precatorius*, Faculty of Medicine, Kynsey Road, Colombo 8, Sri Lanka.
- [5]. Hossain E, Chandra G, Nandy AP, Mandal SC & Gupta JK Anthelmintic effect of a methanol extract of leaves of *Dregea volubilis* on *Paramphistomum explanatum*. *Parasitol Res* 110, 2012, 809–814.
- [6]. Kosalge SB, Fursule RA Investigation of In vitro Anthelmintic activity of *Thespesia Lampas* (Cav.). *Asian J Pharm Clin Res* 2(2), 2009, 69-71.

- [7]. Kokate CK, Purohit AP, Gokhale SB Text Book of Pharmacognosy. 41st ed. Pune: Nirali Prakashani; K.R.Karitikaran; B.D. Basu , Indian medicinal plants, I, 2008, 711
- [8]. Kumar T, Alexander A, Dewangan D, Nagori K. Anthelmintic Activity of the Whole Plant of Bauhinia purpurea (linn.). Asian J Pharm Clin Res 4(3), 2011, 110-111
- [9]. Tripathi KD. Essentials of Medical Pharmacology. 6th Ed. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd. 2008, 808-810.

**How to cite this article:** S. Selvadurai, M.D. Dhana Raju, A. Venkateswara Rao, In-vitro anthelmintic activity of methanolic leaf extract of Abrus precatorius Linn. (fabaceae). Int J of Allied Med Sci and Clin Res 2016; 4(3): 344-348.